



WAHG

WELLINGTON
ANIMAL HOSPITAL
GROUP

WELLINGTON ANIMAL HOSPITAL:

Tel No: 021 873 1196 | **Fax No:** 021 873 3255

32 Piet Retief Street, Wellington
P.O. Box 499, Wellington, 7654

BOULEVARD ANIMAL HOSPITAL - PAARL:

Tel No: 021 863 3187 | **Fax No:** 021 863 2378

72 Berg River Boulevard, Paarl

FRANSCHHOEK ANIMAL CLINIC:

Tel No: 021 876 2504 | **Fax No:** 086 246 7913

Village Artisan, 54 Cabrierg Street, Franschhoek

www.wahg.co.za

Veterinary Report

Patient Name: G2/31

Species: Canine

Breed: Mixed Breed

Age: 4 years

Sex: Female

Presenting Complaint:

Traumatic wound to the left forelimb.

History:

The patient presented with a traumatic injury to the left arm. The injury was sustained recently. There were no other significant medical concerns reported at this time.

Clinical Examination:

Upon physical examination, the left forelimb exhibited a deep laceration with significant tissue damage. The wound showed signs of moderate swelling and mild erythema around the margins. The remainder of the physical examination was unremarkable.

Procedure:

The patient was sedated to facilitate proper examination and wound cleaning. The wound was carefully debrided to remove necrotic tissue and any foreign material. Due to the depth of the injury, a sterile bandage was applied to control swelling and prevent further contamination.

Treatment Plan:

1. **Synulox (Amoxicillin/Clavulanate):** Prescribed to address potential bacterial infection and promote healing.
2. **Meloxicam:** Administered as a non-steroidal anti-inflammatory to manage pain and inflammation.
3. **Bandage:** The bandage will remain in place for 3 days, after which it will be removed to assess wound healing.
4. **Sutures:** Sutures will be removed in 14 days.

Prognosis:

With appropriate care and monitoring, the prognosis for recovery is good. The patient will need to remain under close observation for signs of infection or complications related to the wound healing process.

Follow-up Instructions:

- Keep the bandage clean and dry until the scheduled removal date.
- Administer prescribed medications as directed.
- Return for a recheck in 3 days to evaluate the wound and remove the bandage.
- Return in 14 days for suture removal.
- Monitor for signs of infection, including increased swelling, redness, or discharge from the wound.

wellington@wahg.co.za | boulevard@wahg.co.za | franschhoek@wahg.co.za

Partners: Dr EB Compion, Dr E du Plessis, Dr JB Morkel, Dr GJ van Tonder, Dr PW van Zyl

Reg No: 2015/299686/07 | Vat No: 4680152586